



How many boundary conditions (on the potential V) do you use to find V inside the spherical plastic shell?

- A. 1
- B. 2
- C. 3
- D. 4
- E. It depends on $V_0(\theta)$



ANNOUNCEMENTS


- Homework 8 has 2D relaxation problem
 - It is OK to post code on piazza and get help
 - Solution to HW7 problem 5 (1D relaxation) is linked (you may work from it)
- DC out of town Monday; Norman Birge will cover




Two charges are positioned as shown to the left. The relative position vector between them is \mathbf{d} . What is the value of the dipole moment? $\sum_i q_i \mathbf{r}_i$

- A. $+q\mathbf{d}$
- B. $-q\mathbf{d}$
- C. Zero
- D. None of these

MULTIPOLE EXPANSION



Multipole Expansion of the Power Spectrum of CMBR



Two charges are positioned as shown to the left. The relative position vector between them is \mathbf{d} . What is the dipole moment of this configuration?

$$\sum_i q_i \mathbf{r}_i$$

A. $+q\mathbf{d}$

B. $-q\mathbf{d}$

C. Zero


D. None of these; it's more complicated than before!

For a dipole at the origin pointing in the z-direction, we have derived:


$$\mathbf{E}_{dip}(\mathbf{r}) = \frac{p}{4\pi\epsilon_0 r^3} (2 \cos \theta \hat{\mathbf{r}} + \sin \theta \hat{\boldsymbol{\theta}})$$

For the dipole $\mathbf{p} = q\mathbf{d}$ shown, what does the formula predict for the direction of $\mathbf{E}(\mathbf{r} = 0)$?

- A. Down
- B. Up
- C. Some other direction
- D. The formula doesn't apply




IDEAL VS. REAL DIPOLE



$$\mathbf{p} = \sum_i q_i \mathbf{r}_i$$

What is the magnitude of the dipole moment of this charge distribution?

- A. qd
- B. $2qd$
- C. $3qd$
- D. $4qd$
- E. It's not determined




$$\mathbf{p} = \sum_i q_i \mathbf{r}_i$$

What is the dipole moment of this system?

(BTW, it is NOT overall neutral!)

- A. $q\mathbf{d}$
- B. $2q\mathbf{d}$
- C. $\frac{3}{2}q\mathbf{d}$
- D. $3q\mathbf{d}$
- E. Something else (or not defined)




$$\mathbf{p} = \sum_i q_i \mathbf{r}_i$$

What is the dipole moment of this system?

(Same as last question, just shifted in z .)

- A. $q\mathbf{d}$
- B. $2q\mathbf{d}$
- C. $\frac{3}{2}q\mathbf{d}$
- D. $3q\mathbf{d}$
- E. Something else (or not defined)



You have a physical dipole, $+q$ and $-q$ a finite distance d apart. When can you use the expression:

$$V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \frac{\mathbf{p} \cdot \hat{\mathbf{r}}}{r^2}$$

- A. This is an exact expression everywhere.
- B. It's valid for large r
- C. It's valid for small r
- D. No idea...

You have a physical dipole, $+q$ and $-q$ a finite distance d apart. When can you use the expression:

$$V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \sum_i \frac{q_i}{\mathcal{R}_i}$$

- A. This is an exact expression everywhere.
- B. It's valid for large r
- C. It's valid for small r
- D. No idea...

Which charge distributions below produce a potential that looks like $\frac{C}{r^2}$ when you are far away?

+2q
●

A)

+2q
●
●
+2q

B)

+q +q
● ●
● ●
-q -q

C)



+2q +q
● ●
● ●
-q -2q

D)

E) None of these, or more than one of these!

(For any which you did not select, how DO they behave at large r ?)

Which charge distributions below produce a potential that looks like $\frac{C}{r^2}$ when you are far away?




E) None of these, or more than one of these!

(For any which you did not select, how DO they behave at large r ?)

In terms of the multipole expansion

$V(r) = V(\textit{mono}) + V(\textit{dip}) + V(\textit{quad}) + \dots$, the following charge distribution has the form:



- A. $V(r) = V(\textit{mono}) + V(\textit{dip}) +$ higher order terms
- B. $V(r) = V(\textit{dip}) +$ higher order terms
- C. $V(r) = V(\textit{dip})$
- D. $V(r) =$ only higher order terms than dipole
- E. No higher terms, $V(r) = 0$ for this one.