

Two small spheres (mass, m) are attached to insulating strings (length, L) and hung from the ceiling as shown.

How does the angle (with respect to the vertical) that the string attached to the $-q$ charge (θ_1) compare to that of the $-2q$ charge (θ_2)?

- A. $\theta_1 > \theta_2$
- B. $\theta_1 = \theta_2$
- C. $\theta_1 < \theta_2$
- D. ????

ANNOUNCEMENTS

- CAPS Connect
 - CAPS Connect is a brief consultation program that is confidential, completely free, and available to all enrolled MSU students.
 - Common concerns include: Stress; Difficulty adjusting to school; Academic concerns; Family, roommate, or other relationship issues; Financial concerns; Sadness

Available drop in times

- BPS 1312 - Mondays 9-10:30am


CLASSICAL ELECTROMAGNETISM




$\sim 10^8 \text{ m}$ $\longrightarrow \longrightarrow \longrightarrow \longrightarrow \longrightarrow \longrightarrow \longrightarrow \longrightarrow \sim 10^{-16} \text{ m}$

24 orders of magnitude


1 Å = 100,000 fm



ELECTROSTATICS




5 charges, q , are arranged in a regular pentagon, as shown.
What is the E field at the center?



- A. Zero
- B. Non-zero
- C. Really need trig and a calculator to decide

1 of the 5 charges has been removed, as shown. What's the E field at the center?



A. $+(kq/a^2)\hat{y}$

B. $-(kq/a^2)\hat{y}$

C. 0

D. Something entirely different!

E. This is a nasty problem which I need more time to solve

If all the charges live on a line (1-D), use:

$$\lambda \equiv \frac{\text{charge}}{\text{length}}$$


Draw your own picture. What's $\mathbf{E}(\mathbf{r})$?

To find the E-field at P from a thin line (uniform charge density λ):

$$\mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int \frac{\lambda dl'}{\mathcal{R}^2} \hat{\mathcal{R}}$$

What is \mathcal{R} ?

- A. x
- B. y'
- C. $\sqrt{dl'^2 + x^2}$
- D. $\sqrt{x^2 + y'^2}$
- E. Something else



$$\mathbf{E}(\mathbf{r}) = \int \frac{\lambda dl'}{4\pi\epsilon_0 \mathcal{R}^3} \vec{\mathcal{R}}, \text{ so: } E_x(x, 0, 0) = \frac{\lambda}{4\pi\epsilon_0} \int \dots$$

A. $\int \frac{dy' x}{x^3}$

B. $\int \frac{dy' x}{(x^2 + y'^2)^{3/2}}$

C. $\int \frac{dy' y'}{x^3}$

D. $\int \frac{dy' y'}{(x^2 + y'^2)^{3/2}}$

E. Something else

