


## True or False

"Frozen" or "Baked-In" Polarization means that the dipoles in the material will shift when an external field is present.

- A. True
- B. False
- C. ???

A solid non-conducting dielectric rod has been injected ("doped") with a fixed, known charge distribution  $\rho(s)$ . (The material responds, polarizing internally.)



When computing  $D$  in the rod, do you treat this  $\rho(s)$  as the "free charges" or "bound charges"?

- A. "free charge"
- B. "bound charge"
- C. Neither of these -  $\rho(s)$  is some combination of free and bound
- D. Something else.

We define "Electric Displacement" or "D" field,

$$\mathbf{D} = \epsilon_0 \mathbf{E} + \mathbf{P}$$


If you put a dielectric in an **external** field, it polarizes, adding a new **induced** field (from the bound charges). These superpose, making a **total** electric field. Which of these three E fields is the "E" in the formula for D above?

- A.  $\mathbf{E}_{ext}$
- B.  $\mathbf{E}_{induced}$
- C.  $\mathbf{E}_{tot}$

We define  $\mathbf{D} = \epsilon_0 \mathbf{E} + \mathbf{P}$ , with

$$\oint \mathbf{D} \cdot d\mathbf{A} = Q_{free}$$

A point charge  $+q$  is placed at the center of a dielectric sphere (radius  $R$ ). There are no other free charges anywhere. What is  $|\mathbf{D}(r)|$ ?



- A.  $q/(4\pi r^2)$  everywhere
- B.  $q/(4\epsilon_0\pi r^2)$  everywhere
- C.  $q/(4\pi r^2)$  for  $r < R$ , but  $q/(4\epsilon_0\pi r^2)$  for  $r > R$
- D. None of the above, it's more complicated
- E. We need more info to answer!

For linear dielectrics the relationship between the polarization,  $\mathbf{P}$ , and the total electric field,  $\mathbf{E}$ , is given by:

$$\mathbf{P} = \epsilon_0 \chi_e \mathbf{E}$$

where  $\chi_e$  is typically a known constant. Think about what happens if (1)  $\chi_e \rightarrow 0$  or if (2)  $\chi_e \rightarrow \infty$ . What do each of these limits describe?


- A. (1) describes a metal and (2) describes vacuum
- B. (1) describes vacuum and (2) describes a metal
- C. Any material can give either  $\chi_e \rightarrow 0$  or  $\chi_e \rightarrow \infty$

When there are no free charges,  $\rho_{free} = 0$ , in a linear dielectric material, the electric potential,  $V$ , in that material satisfies Laplace's equation.

$$\nabla^2 V = 0$$

- A. True
- B. False
- C. ???

A very large (effectively infinite) capacitor has charge  $Q$ . A neutral (*homogeneous*) dielectric is inserted into the gap (and of course, it will polarize). We want to find  $\mathbf{E}$  everywhere.



Which equation would you head to first?

A.  $\mathbf{D} = \epsilon_0 \mathbf{E} + \mathbf{P}$


B.  $\oint \mathbf{D} \cdot d\mathbf{A} = Q_{free}$

C.  $\oint \mathbf{E} \cdot d\mathbf{A} = \frac{Q}{\epsilon_0}$

D. More than one of these would work

E. Can't solve unless we know the dielectric is linear.

A very large (effectively infinite) capacitor has charge  $Q$ . A neutral (*homogeneous*) dielectric is inserted into the gap (and of course, it will polarize). We want to find  $\mathbf{D}$  everywhere.



Which equation would you head to first?

A.  $\mathbf{D} = \epsilon_0 \mathbf{E} + \mathbf{P}$


B.  $\oint \mathbf{D} \cdot d\mathbf{A} = Q_{free}$

C.  $\oint \mathbf{E} \cdot d\mathbf{A} = \frac{Q}{\epsilon_0}$

D. More than one of these would work



An ideal (large) capacitor has charge  $Q$ . A neutral linear dielectric is inserted into the gap. We want to find  $\mathbf{D}$  in the dielectric.




$$\oint \mathbf{D} \cdot d\mathbf{A} = Q_{free}$$

For the Gaussian pillbox shown, what is  $Q_{free,enclosed}$ ?

- A.  $\sigma A$
- B.  $-\sigma_B A$
- C.  $(\sigma - \sigma_B) A$
- D.  $(\sigma + \sigma_B) A$
- E. Something else

An ideal (large) capacitor has charge  $Q$ . A neutral linear dielectric is inserted into the gap. We want to find  $\mathbf{D}$  in the dielectric.




$$\oint \mathbf{D} \cdot d\mathbf{A} = Q_{free}$$

Is  $\mathbf{D}$  zero INSIDE the metal? (i.e., on the top face of our cubical Gaussian surface)

- A. It must be zero in there.
- B. It depends.
- C. It is definitely not zero in there.

An ideal (large) capacitor has charge  $Q$ . A neutral linear dielectric is inserted into the gap. We want to find  $\mathbf{D}$  in the dielectric.



$$\oint \mathbf{D} \cdot d\mathbf{A} = Q_{free}$$

What is  $|\mathbf{D}|$  in the dielectric?

- A.  $\sigma$
- B.  $2\sigma$
- C.  $\sigma/2$
- D.  $\sigma + \sigma_b$
- E. Something else