

The electric fields of two EM waves in vacuum are both described by:

$$\mathbf{E} = E_0 \sin(kx - \omega t)\hat{y}$$

The "wave number"  $k$  of wave 1 is larger than that of wave 2,  $k_1 > k_2$ . Which wave has the larger frequency  $f$ ?

- A. Wave 1
- B. Wave 2
- C. impossible to tell

For a wave on a 1d string that hits a boundary between 2 strings of different material we get,

$$\begin{aligned}\tilde{f}(z < 0) &= \tilde{A}_I e^{i(k_1)z - \omega t} + \tilde{A}_R e^{i(-k_1)z - \omega t} \\ \tilde{f}(z > 0) &= \tilde{A}_T e^{i(k_2)z - \omega t}\end{aligned}$$

where continuity (BCs) give,

$$\begin{aligned}\tilde{A}_R &= \left( \frac{k_1 - k_2}{k_1 + k_2} \right) \tilde{A}_I \\ \tilde{A}_T &= \left( \frac{2k_1}{k_1 + k_2} \right) \tilde{A}_I\end{aligned}$$

Is the transmitted wave in phase with the incident wave?

- A) Yes, always B) No, never C) Depends

For a wave on a 1d string that hits a boundary between 2 strings of different material we get,

$$\begin{aligned}\tilde{f}(z < 0) &= \tilde{A}_I e^{i(k_1)z - \omega t} + \tilde{A}_R e^{i(-k_1)z - \omega t} \\ \tilde{f}(z > 0) &= \tilde{A}_T e^{i(k_2)z - \omega t}\end{aligned}$$

where continuity (BCs) give,


$$\begin{aligned}\tilde{A}_R &= \left( \frac{k_1 - k_2}{k_1 + k_2} \right) \tilde{A}_I \\ \tilde{A}_T &= \left( \frac{2k_1}{k_1 + k_2} \right) \tilde{A}_I\end{aligned}$$

Is the reflected wave in phase with the incident wave?

- A) Yes, always B) No, never C) Depends

An electromagnetic plane wave propagates to the right. Four vertical antennas are labeled 1-4. 1, 2, and 3 lie in the  $x - y$  plane. 1, 2, and 4 have the same  $x$ -coordinate, but antenna 4 is located further out in the  $z$ -direction. Rank the time-averaged signals received by each antenna.

- A.  $1=2=3>4$
- B.  $3>2>1=4$
- C.  $1=2=4>3$
- D.  $1=2=3=4$
- E.  $3>1=2=4$



A point source of radiation emits power  $P_0$  isotropically (uniformly in all directions). A detector of area  $a_d$  is located a distance  $R$  away from the source. What is the power  $P_d$  received by the detector?

- A.  $\frac{P_0}{4\pi R^2} a_d$
- B.  $P_0 \frac{a_d^2}{R^2}$
- C.  $P_0 \frac{a_d}{R}$
- D.  $\frac{P_0}{\pi R^2} a_d$
- E. None of these

