


ANNOUNCEMENTS

- Homework 10 (it's long; you started it, right?)
 - Due this Friday
- Final Homework is due Friday the 9th
 - Magnetic dipoles and some magnetic matter
- Final Exam (20%)
 - 12:45pm-2:45pm on Thursday the 15th in this room
- Detailed grade projections by Monday 12th
 - w/ clicker bonus, but not HW 11
- SIRS are open
 - Please fill out; it helps shape departmental offerings

The vector potential \mathbf{A} due to a long straight wire with current I along the z -axis is in the direction parallel to:

- A. \hat{z}
- B. $\hat{\phi}$ (azimuthal)
- C. \hat{s} (radial)

Assume the Coulomb Gauge



Consider a fat wire with radius a with uniform current I_0 that runs along the $+z$ -axis. We can compute the vector potential due to this wire directly. What is \mathbf{J} ?

- A. $I_0/(2\pi)$
- B. $I_0/(\pi a^2)$
- C. $I_0/(2\pi a)\hat{z}$
- D. $I_0/(\pi a^2)\hat{z}$
- E. Something else!?

Consider a fat wire with radius a with uniform current I_0 that runs along the $+z$ -axis. Given $\mathbf{A}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int \frac{\mathbf{J}(\mathbf{r}')}{\mathcal{R}} d\tau'$, which components of \mathbf{A} need to be computed?

- A. All of them
- B. Just A_x
- C. Just A_y
- D. Just A_z
- E. Some combination

Consider line of charge with uniform charge density, $\lambda = \rho\pi a^2$. What is the magnitude of the electric field outside of the line charge (at a distance $s > a$)?

- A. $E = \lambda/(4\pi\epsilon_0 s^2)$
- B. $E = \lambda/(2\pi\epsilon_0 s^2)$
- C. $E = \lambda/(4\pi\epsilon_0 s)$
- D. $E = \lambda/(2\pi\epsilon_0 s)$
- E. Something else?!

Use Gauss' Law

Consider a shell of charge with surface charge σ that is rotating at angular frequency of ω . Which of the expressions below describe the surface current, \mathbf{K} , that is observed in the fixed frame.

- A. $\sigma \omega$
- B. $\sigma \dot{\mathbf{r}}$
- C. $\sigma \mathbf{r} \times \omega$
- D. $\sigma \omega \times \mathbf{r}$
- E. Something else?